

Research Article

IN-OVO TREATMENT OF CHLORPYRIFOS AND CYPERMETHRIN IN COMBINATION ALTERED THE HAEMATOLOGICAL PARAMETERS IN TWO GENERATIONS OF DOMESTIC HEN

Asiya N. Khan, Gowri K. Uggini, Sonam P. Patel and *Suresh Balakrishnan

Department of Zoology, Faculty of Science, the Maharaja Sayajirao University of Baroda, Vadodara – 390002, Gujarat, India

** Author for Correspondence*

ABSTRACT

Effects of in-ovo intoxication of an insecticide formulation Nurelle D 505 EC (chlorpyrifos 50% EC + cypermethrin 5% EC) was evaluated in two generations of domestic fowl. The investigation covered four groups of fertilized RIR eggs - three for experimental and one for control. Experimental groups of eggs received doses of 0.01, 0.05 and 0.1 µg/egg of combination insecticide (Ci) and the control received corn oil on day zero of incubation. The doses were selected based on our previous study. The eggs were incubated and upon hatching the F1 generation chicks were sorted as per their treatment, tagged and reared on standard diet. Upon sexual maturity the birds from the same groups were allowed pen mating. The fertilized eggs were collected and incubated for the next generation (F2). Blood samples were drawn from the brachial vein of 25 weeks old parent F1 generation and 4 weeks old chick of F2 generation and standard haematological parameters like TRBC, TWBC, DC, Hb as well as PCV were recorded and hemogramme was computed. The study showed that chlorpyrifos and cypermethrin applied in a mixture caused significant reduction in the RBC, PCV, lymphocytes and platelets in both F1 and F2 generations, while the WBC count was lowered in F1 and raised in F2 generation. The apparent erythropenia condition observed in the F2 generation of Ci treated chicks could be attributed to a compromised erythropoiesis rather than direct cytotoxicity and internal haemorrhage though the molecular basis of the same remains to be identified. Further, lymphocytopenia an indicator of reduced immunocompetence was also observed in chicks of F2 generation. This is the first report of Ci induced haemetological alteration in the second generation of chicks and we believe that the test system can evolve as an alternate model for testing two generation developmental toxicity.

Keywords: *Pesticides, Chlorpyrifos, Cypermethrin, Haematology, Fertilized RIR Eggs*

INTRODUCTION

Pest control measures are used world over to protect largely the agro-ecosystem (Bhattacharya and Bhattacharya, 2006). In mid 1960s with agricultural modernization in developing countries, the global pesticide industry witnessed a prodigious growth (Binswanger *et al.*, 1998). Sporadic use of pesticides has now become prevalent and has led to an extensive contamination of the environment. The research analysis shows that annual consumption of pesticides is about 24% in United States, 45% in Europe, and 31% in the rest of the world (Foo and Hameed, 2010). However, because of injudicious use of pesticides in India, 51% of food commodities are contaminated with pesticide residues (Gupta, 2004). The predominant classes of pesticides that are in use include organophosphates, carbamates and pyrethroids (Masri *et al.*, 2009).

Chlorpyrifos is one of the most widely used organophosphorous insecticide and belongs to class II of World Health Organization classification of pesticides (Cox, 1997). Chlorpyrifos inhibits acetylcholine decomposition and hence increases the acetylcholine level in the synaptic cleft and stimulates specific receptors (Nostrandt *et al.*, 1997). Chlorpyrifos is moderately toxic and a prolonged exposure has been reported to cause anemia although the mechanism has not been elucidated (Ambali *et al.*, 2010).

Cypermethrin is an active pyrethroid which is known to stimulate nerves by causing pronounced repetitive activity. It has been reported that synapses are more sensitive to the pyrethroids than nerve fibres (Latuszynska *et al.*, 2001). Mechanistically, pyrethroid binds to and delay sodium channel closure,

Research Article

so that a prolonged sodium tail current persists after the membrane repolarization (Perger and Szadkowschi, 1994). Nevertheless, this neuro-impairment might result in other physiological manifestations, which can be gauged through haemogram analysis.

In the present study, a commercially available combination insecticide (chlorpyrifos + cypermethrin) (Ci) was used to analyze the haematological changes in the first filial (F1) and second filial (F2) generations of domestic fowls.

The rationale behind the selection of the present test article is that, chlorpyrifos and cypermethrin are amongst the most frequently used insecticides. Further, due to the potentiation of the effects of cypermethrin by organophosphorus compound, has resulted in the promotion of combination insecticide on a commercial scale. Therefore it was thought pertinent to analyze the toxic effect of Ci rather than individual pesticide.

It is well known that haematological values indicate the state of health and has been used as a diagnostic tool by the clinicians for effective treatment. Hence, the objective of present study was to evaluate the two generation developmental toxicity potential of the chlorpyrifos and cypermethrin combination in domestic fowl through complete blood cell count of both F1 and F2 generation.

MATERIALS AND METHODS

Test Chemical

Commercially available insecticide Nurelle D 505 EC (chlorpyrifos 50% EC + cypermethrin 5% EC, Dow AgroSciences, USA) was used in the study.

Egg Procurement

The fertilized Rhode Island-Red (RIR) eggs were collected from Poultry Science Division of Anand Agricultural University, Anand, and refrigerated at 4°C until used. Eggs were swabbed with povidone-iodine to avoid infection. The investigation covered four groups of fifty fertilized RIR eggs each; three for experimental and one control. Each egg was marked to receive the respective dosage treatment of the Ci for experimental groups and corn oil for control group. The study was conducted on the F1 (Parents) and F2 (Chicks) generations as the animal model.

Insecticide Injection

Combination insecticide was diluted in corn oil and three sub-lethal doses namely low, mid and high were made as described in our earlier studies (Uggini *et al.*, 2012). A single dose of 0.01, 0.05 and 0.1 µg/egg of Ci diluted in corn oil was administered to the airspaces of fertilized eggs from low, mid and high group respectively on day zero of incubation. The eggs from the control received corn oil in the similar manner. The injected portion of each egg was sealed by molten paraffin wax. The whole experiment was done in a sterile environment under a laminar air flow. The dose volume of Ci and corn oil was maintained at 25µl/egg.

Incubation

A cleaned, disinfected and fumigated automatic egg incubator and hatcher were used. Incubator regulates the factors such as temperature of $37.5 \pm 0.5^{\circ}\text{C}$, 75-80% relative humidity and periodic turning of the eggs. Immediately after treatment, both the treated and control eggs were kept in the incubator with proper marking for 18 days. On 18th day the egg candling was done and the viable eggs were transferred to hatcher till the day of hatch. On day 21 after hatching, the hatchling groups (F1 generation) from the treated and control eggs were tagged with wing bands with respect to their dosage treatment and were housed in separate pens with immediate supply of water. After three hours starter mash feed was provided to the chicks. To promote the survival of the chick, assisted hatching was involved in case of those chicks who failed to progress normally at hatching stage due to deformities or weakness caused by Ci.

The F1 generation chicks when reached 25 weeks old were randomized within the respective treatment groups and the similar weighed ones were selected for breeding. Twenty hens and two roosters from each of the four groups were selected and pen mated for a week. The eggs collected from the F1 generation were marked as per the treatment groups of F1 parental generation and incubated. On hatch out the F2 generation chicks were wing banded with respect to their parent's group and reared on standard diet for 4

Research Article

weeks. The haematological analyses were performed on 25 week old parents and 4 week old second generation chicks.

Haematological Estimations

Blood samples were drawn from brachial vein of 25 week old parent and 4 week old F2 chick using EDTA rinsed 2ml disposable syringes and collected into EDTA rinsed or heparinized vials. After collection, the samples were refrigerated and processed within 6 hr.

Hematology analyzer, Mindray BC2300 (Shenzhen Mindray Bio-Medical Electronics Co., Ltd., China) was used to evaluate haematological parameters including total erythrocyte (RBC) count, haemoglobin (Hb) concentration, packed cell volume (PCV), total leukocyte (WBC) count and differential leukocyte counts. Complementary indices of the erythrocytic component, including mean corpuscular haemoglobin concentration - MCHC, mean corpuscular haemoglobin - MCH, and mean corpuscular volume - MCV, were also computed.

Effect of treatment on differential white blood cell counts was enumerated in blood smears prepared and stained panoptically with Microkrom Microscopy Stains (Giemsa Stain) immediately after sampling.

Raw data were processed and analyzed by one-way ANOVA followed by Dunnet's multiple comparison test. Results were expressed as mean \pm standard deviation. A 'p' value less than or equal to 0.05 was considered as statistically significant. Statistical analysis was performed using GraphPad Prism 5. The protocol mentioned herein was approved by the IAEC of the department in full compliance with the CPCSEA norms.

RESULTS AND DISCUSSION

Results

The purpose of the present study was to evaluate the possibility of haematologic abnormalities, if any, in the F2 generation induced by Ci treatment to the eggs that gave rise to F1 generation. The haematological parameters in F1 and F2 generations of chicks were analyzed after administering a single dose of 0.01, 0.05 and 0.10 $\mu\text{g/egg}$ of Ci to the air cell of eggs and apparent variations in the haematological parameters with that of a control group were recorded.

F1 Generation

At 0.01 $\mu\text{g/egg}$ and 0.05 $\mu\text{g/egg}$ of Ci, the F1 generation did not show any significant changes in haemoglobin content when compared to the concurrent controls whereas treatment with 0.1 $\mu\text{g/egg}$ of Ci induced severe anaemia ($p \leq 0.05$) in the first generation of chicks.

Total red blood cell concentration values were recorded for the birds that received low dose (0.01 $\mu\text{g/egg}$) of Ci, and were found comparable to that of controls. The analysis of blood showed a decline in the total number of red blood cells at 0.05 $\mu\text{g/egg}$ and 0.10 $\mu\text{g/egg}$ of Ci as compared to the control group. The PCV value also showed significant reduction in the group that received 0.1 $\mu\text{g/egg}$. However, the MCV values showed an apparent hike in F1 generation while MCH values were found decreased in all the three treatment groups compared to control group. Though MCHC value showed a similar declining trend, it was found statistically non significant. Quite similar to the trend showed by erythrocytes, the WBC count too was found reduced in birds that received Ci. Nevertheless, the differential white blood cell count showed an increase in the number of polymorphs only in the birds from high dose group, whereas the lymphocyte numbers in this group of birds were found significantly lowered with respect to that of controls. The eosinophil showed an increase in mean number in birds from the F1 groups that received Ci, whereas monocyte number declined in all the three first generation groups of chicks. Basophil did not show any significant change in its number at any of the selected doses. The platelet count however, recorded significant reduction in count in birds that received high dose of Ci (**Table 1**).

F2 Generation

The haemoglobin content has not shown any significant variation in any group of the F2 generation chicks that descended from a parent who received Ci during development. However, the total RBC count was lowered in the groups whose parents received 0.05 and 0.10 $\mu\text{g/egg}$ of Ci. The other parameters like PCV, MCV, MCH, MCHC count have not shown any significant alterations in any of the F2 generation

Research Article

chicks. However, birds from the F2 generation of the high dose group showed an increase in the mean WBC count. The differential count of the white blood cells showed that the level of polymorphs was significantly increased in all the Ci treated groups compared to the control. Nonetheless, the lymphocyte eosinophil, monocytes and platelet counts were found significantly decreased in all the three F2 generation chicks descended from Ci treated parents. However, basophil did not show any significant treatment related changes in any of the treated animals with respect to the control group animals (Table 2).

Table 1: Complete blood count of F1 generation RIR domestic fowl subjected to various doses of combination insecticide during their embryonic development.

Parameters	Control (Corn Oil)	Low (0.01ug/egg)	Medium (0.05ug/egg)	High (0.1ug/egg)
Haemoglobin (gm/dl)	10.13 ± 0.79 ^a	9.47 ± 0.91 ^a	9.03 ± 0.85 ^a	5.93 ± 0.57 ^b
RBC Count (10 ⁶ /cumm)	3.77 ± 0.24 ^a	3.70 ± 0.35 ^a	3.09 ± 0.31 ^b	2.84 ± 0.28 ^b
P.C.V (%)	30.10 ± 2.71 ^a	29.72 ± 2.84 ^a	27.02 ± 2.70 ^a	18.65 ± 1.79 ^b
M.C.V (fl)	80.72 ± 7.89 ^a	85.96 ± 8.42 ^a	88.57 ± 8.74 ^a	97.68 ± 9.62 ^b
M.C.H (pg)	27.14 ± 2.43 ^a	25.71 ± 2.57 ^b	23.14 ± 2.38 ^c	20.27 ± 1.98 ^c
M.C.H.C (g/dl)	33.68 ± 2.33 ^a	33.30 ± 2.58 ^a	32.86 ± 3.10 ^a	31.94 ± 2.99 ^a
WBC Count (/cumm)	13450 ± 505.77 ^a	11883 ± 527.65 ^b	10700 ± 532.82 ^b	10400 ± 530.09 ^b
Polymorphs (%)	57.50 ± 3.51 ^a	64.67 ± 4.33 ^a	66.33 ± 5.97 ^a	67.33 ± 6.73 ^b
Lymphocytes (%)	38.00 ± 2.12 ^a	29.50 ± 2.27 ^b	28.33 ± 2.80 ^c	25.33 ± 2.12 ^d
Eosinophils (%)	2.17 ± 0.19 ^a	3.17 ± 0.29 ^b	3.33 ± 0.32 ^b	3.83 ± 0.39 ^c
Monocytes (%)	2.33 ± 0.17 ^a	2.00 ± 0.21 ^b	1.50 ± 0.14 ^c	1.33 ± 0.11 ^c
Basophils (%)	1.02 ± 0.33 ^a	2.05 ± 0.70 ^a	1.50 ± 0.72 ^a	2.04 ± 0.65 ^a
Platelet Count (K/cumm)	2.33 ± 0.14 ^a	2.26 ± 0.19 ^a	2.14 ± 0.20 ^a	1.79 ± 0.16 ^b

Table 2: Complete blood count of F2 generation chicks descended from parents who received low, medium or high doses of Ci during their embryonic development.

Parameters	Control (Corn Oil)	Low (0.01ug/egg)	Medium (0.05ug/egg)	High (0.1ug/egg)
Haemoglobin (gm/dl)	10.19 ± 0.85 ^a	10.03 ± 0.92 ^a	9.80 ± 0.91 ^a	9.42 ± 0.90 ^a
RBC Count (10 ⁶ /cumm)	4.16 ± 0.31 ^a	3.97 ± 0.33 ^a	2.74 ± 0.27 ^b	2.32 ± 0.25 ^b
P.C.V (%)	30.51 ± 3.03 ^a	29.92 ± 2.94 ^a	26.77 ± 2.69 ^a	26.32 ± 2.55 ^a
M.C.V (fl)	78.83 ± 7.56 ^a	79.50 ± 7.82 ^a	85.99 ± 8.49 ^a	87.84 ± 8.73 ^a
M.C.H (pg)	26.34 ± 2.33 ^a	24.95 ± 2.41 ^a	24.26 ± 2.42 ^a	23.20 ± 2.34 ^a
M.C.H.C (g/dl)	33.38 ± 2.98 ^a	33.15 ± 3.24 ^a	32.59 ± 3.19 ^a	32.22 ± 3.16 ^a
WBC Count (/cumm)	14057 ± 578.32 ^a	14116 ± 620.30 ^a	14533 ± 655.67 ^a	15150 ± 724.47 ^b
Polymorphs (%)	27.14 ± 2.13 ^a	56.50 ± 3.78 ^b	59.67 ± 4.66 ^b	61.83 ± 5.77 ^b
Lymphocytes (%)	68.00 ± 3.21 ^a	39.33 ± 2.39 ^a	34.50 ± 3.22 ^a	34.33 ± 3.01 ^a
Eosinophils (%)	2.57 ± 0.17 ^a	2.17 ± 0.21 ^b	1.83 ± 0.18 ^c	1.67 ± 0.16 ^c
Monocytes (%)	2.29 ± 0.19 ^a	2.00 ± 0.89 ^b	1.50 ± 0.14 ^b	1.33 ± 0.11 ^b
Basophils (%)	1.71 ± 0.42 ^a	1.94 ± 0.62 ^a	1.34 ± 0.55 ^a	2.12 ± 0.64 ^a
Platelet Count (K/cumm)	3.16 ± 0.23 ^a	2.55 ± 0.21 ^b	2.24 ± 0.22 ^b	2.11 ± 0.20 ^c

Values are expressed in mean ± SD; Values with same superscript are not statistically significant for each parameter.

RBC-Red blood corpuscles, WBC-White blood corpuscles, P.C.V.-Packed cell volume, M.C.V.-Mean corpuscular volume, M.C.H.-Mean corpuscular haemoglobin, M.C.H.C-Mean corpuscular haemoglobin concentration.

Research Article

Discussion

A scan through the literature has revealed that the pesticide intoxication often results in altered hemogramme. According to earlier reports, variations in the haematological parameters were observed as a result of cypermethrin toxicity (Morgan *et al.*, 1980) and chlorpyrifos toxicity (Barna-Lloyd *et al.*, 1991; Goel *et al.*, 2006; Ambali, 2009) where they reported a decrease in RBC count, Hb and PCV levels. Tung *et al.*, 1975, opined that these changes could be due to the destruction of RBCs with integral disability, which are formed by a hyperactive bone marrow. Moreover, the decrease in RBC counts is reported to be symptomatic of erythrocyte damage or erythrocyte formation (Shakoori *et al.*, 1990). It has been documented that the poisoning by pesticide residues results in anaemia due to interference of haemoglobin biosynthesis and shortening of the lifespan of circulating erythrocytes (Betrosian *et al.*, 1995; Jyostana *et al.*, 2003). Furthermore, cypermethrin is known to get accumulated in cell membranes and hence, disturb the membrane integrity which ultimately leads to the lysis of erythrocytes resulting in erythrocytopenia (Michelangeli *et al.*, 1990). The low Hb concentration observed in the chicks from the Ci treated group may be attributed to the enhanced Hb destruction or declined Hb formation as opined by several earlier workers (Moss and Hathway, 1964, Vural *et al.*, 1986). However, hyperactivity of bone marrow or rupture and destruction of red blood cells might also contribute to the induction of anaemia in case of pesticide poisoning (Barger, 2003).

An increase in MCV and decrease in MCH was observed in all the treated groups of F1 generation compared to their control group indicating macrocytic and hypochromic anemia. Barger, (2003), while explaining the reasons for macrocytic and hypochromic anemia, cited hyper activity of bone marrow or deficiency of some haemopoietic factors as the possible causative factors and a same can be applied to explain the pathology observed in the current study as well. The increase in MCV indicates presence of abnormally large sized erythrocytes (Latimer *et al.*, 2004). Increased MCV may also be observed in regenerative anemia due to hemolysis and haemorrhages. However, the F2 generation has not shown any significant changes in Hb, MCV and MCH, though there was a reduction in the total RBC count. Generally, a decline in total red blood cell counts along with a decrement in PCV, while the MCV, MCH and MCHC being normal, indicates that the anaemic condition is due to hampered erythropoiesis. Though direct cytotoxicity and/or internal haemorrhages inflicted by the insecticide might also result in erythrocytopenia, their contribution in the present case of F2 generation is least likely. There are reports that when internal haemorrhages occur, the erythrocytes are absorbed by the lymphatic vessels or are lysed or phagocytosed (Latimer *et al.*, 2004) and hence lead to a reduced RBC count.

Here in this study leucopenia has been recorded in F1 generation the same alteration was previously reported in chicks treated with the pyrethroid compound fenvalerate (Garg *et al.*, 2004) and the organophosphates chlorpyrifos and methidathion (Ojezele and Abatan, 2009). Further, the decrease in total white blood cell count and lymphocytes in 0.05 and 0.1µg/egg of Ci treated groups may indicate adversely affected lymphoid progenitors. The heterophilia observed in 0.1µg/egg of Ci treated group, might be an immune response to a metabolic or chemical poisoning and /or tissue necrosis. Similar observations were made by Cho *et al.*, (1989), who found lymphocytopenia and neutrophilia in rats exposed to subchronic doses of organophosphate. It is well known that the number of circulating lymphocytes in peripheral blood is an index of functional ability of lymphoid organs and the lymphocytes are the main agents of immunogenic response in birds as well as mammals. Thus, a reduction in the number of total lymphocyte may be an indication of compromised immunocompetence of the intoxicated birds. In the F2 generation, though the total WBC counts seemed unaltered, heterophilia and lymphocytopenia were observed, which vividly depicts subdued immunocompetence. Gowri and coworkers (2010) have reported that Ci treatment induces immunotoxicity in chicks of F1 generation. This report consolidates the present notion of reduced immunocompetence even in second generation as evidenced by the lymphocytopenia.

Further in the present study, noteworthy decline in platelets are seen in chicks from all the three treatment groups which can be attributed to the Ci induced compromised redox physiology. Araujo and colleagues (2008) have reported oxidative stress as a possible mediator of thrombocytopenia in malaria patients. This

Research Article

report gives credence to our notion of reactive oxygen species as a possible reason for thrombocytopenia in Ci intoxicated chicks. Finally, since all the selected doses induced toxicity to the chicks in a dose dependent manner it could be postulated that NOAEL dose should be less than 0.01 µg/egg.

Conclusion

The study reveals the susceptibility of external toxic substances to not only to the adult animals but also to the descendents. The toxicants influenced the health of parent generation when dosed during their in-ovo development to a great extent and also hampered the embryonic development of the next generation. The haematological analysis divulged that the F2 generation animals had compromised immunity compared to the animals receiving the pesticide dose directly. This finding underlines the need for screening of the known developmental toxicants for more than one generation to garner the much needed evidences behind many mysterious human diseases about which the aetiology is still at large.

ACKNOWLEDGEMENT

The authors are thankful to authorities at the Anand Agricultural University for infrastructural facility. One of the authors SPP is thankful to UGC-BSR for financial support in the form of RFSMS.

REFERENCES

- Al Masri MG, Rasheed M and Alawi M (2009).** Degradation of organochlorine pesticides in carbonate sediments from Aqaba Gulf, Red Sea. *Research Journal of Environmental Toxicology* **3**(1) 47-158.
- Ambali SF (2009).** Ameliorative effect of vitamins C and E on neurotoxicological, hematological and biochemical changes induced by chronic chlorpyrifos in Wistar rats. Ph.D. Thesis, Ahmadu Bello University, Zaria.
- Ambali SF, Abubakar AT, Shittu M, Yaqub LS, Anafi SB and Abdullahi A (2010).** Chlorpyrifos-induced alteration of hematological parameters in Wistar rats: ameliorative effect of zinc. *Research Journal of Environmental Toxicology* **4**(2) 55–66.
- Araujo CF, Lacerda MVG, Abdalla DSP and Lima ES (2008).** The role of platelet and plasma markers of antioxidant status and oxidative stress in thrombocytopenia among patients with vivax malaria. *Memorias do Instituto Oswaldo Cruz* **103**(6) 517–521.
- Barger AM (2003).** The complete blood cell count: a powerful diagnostic tool. *Veterinary Clinics: Small Animal Practice* **33** 1207-1222.
- Barna-Lloyd T, Szabo JR and Davis NL (1991).** Chlorpyrifos-Methyl (Reldan R) Rat Chronic Dietary Toxicity/Oncogenicity Study. WHO, Indianapolis, USA.
- Betrosian A, Balla M, Kafiri G, Kofinas G, Makri R and Kakouri A (1995).** Multiple systems organ failure from organophosphate poisoning. *Journal of Toxicology, Clinical Toxicology* **33**(3) 257-260.
- Bhattacharya S and Bhattacharya DK (2006).** Pest control through viral disease: Mathematical modeling and analysis. *Journal of Theoretical Biology* **238** 177-197.
- Binswanger H, Hazell P and McCalla A (1998).** Agriculture and the Environment: Perspectives on Sustainable Rural Development. International Bank for Reconstruction and Development, World Bank, Washington DC.
- Cho JH, Park JM and Jaen YH (1989).** Studies on the diagnosis of organophosphate insecticides poisoning: Clinical and haematological diagnosis. *Research Reports of the Rural Development Administration* **31** 8-13.
- Cox C (1997).** Fungicide fact sheet: Chlorothalonil. *Journal of Pesticide Reform* **17** 14-20.
- Foo KY and Hameed BH (2010).** Detoxification of pesticide waste via activated carbon adsorption process: Review. *Journal of Hazardous Materials* **175** 1-11.
- Garg UK, Pal AK, Jha GJ and Jadhao SB (2004).** Pathophysiological effects of chronic toxicity with synthetic pyrethroid, organophosphate and chlorinated pesticides on bone health of broiler chicks. *Toxicologic Pathology* **32** 364-369.
- Goel A, Dani V and Dhawan DK (2006).** Role of zinc in mitigating the toxic effects of chlorpyrifos on hematological alterations and electron microscopic observations in rat blood. *Biometals* **19**(5) 483-492.

Research Article

Gowri UK, Patel PV and Suresh B (2010). Evaluation of pesticide induced developmental immunotoxicity in RIR chicks. *Journal of Cell and Tissue Research* **10**(2) 2229-2234.

Gupta PK (2004). Pesticide exposure-Indian scene. *Toxicology* **198** 83-90.

Jyostana AP, Arun JP and Sanjay PG (2003). Biochemical effects of various on sprayersof grape gardens. *Indian Journal of Clinical Biochemistry* **18**(2) 16-22.

Latimer KS, Mahaffey EA and Prasse KW (2004). *Clinical Pathology: Veterinary Laboratory Medicine*, 4th edition (Iowa State Univ. Press, Ames, Iowa, USA).

Latuszynska J, Luty S, Raszewski G, Tokarska –Rodak M, Przebirowska D, Przylepa E and Haratym-Maj A (2001). Neurotoxic effect of dermally-applied chlorpyrifos and cypermethrin in wistar rats. *Annals of Agricultural and Environmental Medicine* **8** 163-170.

Michelangeli F, Robson MJ, East JM and Lee AG (1990). The conformation of pyrethroids bound to lipid layers. *Biochimica et Biophysica Acta* **1028** 49- 57.

Morgan DP, Stockdale EM, Roberts RJ and Walter HW (1980). Anaemia associated with exposure to lindane. *Archives of Environmental Health* **35** 307- 310.

Moss JA and Hathway DE (1964). Transport of organic compounds in the mammalian partition of dieldrin and telodrin between the cellular components and soluble proteins of blood. *Biochemical Journal* **91** 383-93.

Nostrandt AC, Padilla S and Moser VC (1997). The relationship of oral chlorpyrifos effects on behavior, cholinesterase inhibition and muscarine receptor density in rat. *Pharmacology Biochemistry and Behavior* **58** 15-23.

Ojezele MO and Abatan OM (2009). Toxicological effects of chlorpyrifos and methidathion in young chickens. *African Journal of Biochemistry Research* **3** 48-51.

Perger G and Szadkowski D (1994). Toxicology of pyrethroids and their relevance to human health. *Annals of Agricultural and Environmental Medicine* **1** 11-17.

Shakoori AR, Aziz F, Alam J and Ali SS (1990). Toxic effects of talastar, a new synthetic pyrethroid, on blood and liver of rabbit. *Pakistan Journal of Zoology* **23** 289-300.

Tung HT, Cook FW, Wyatt RD and Hamilton PB (1975). The anaemia caused by aflatoxin. *Poultry Science* **54** 1962-1969.

Uggini GK, Patel PV and Balakrishnan S (2012). Embryotoxic and teratogenic effects of pesticides in chick embryos: A comparative study using two commercial formulations. *Environmental Toxicology* **27**(3) 166-174.

Vural S, Cetin ET and Tuzlaci U (1986). *Klinik Teşhiste Laboratuvar*, Nurettin Uycan Cilt ve Basım Sanayii, İstanbul. (In Turkish)