

Research Article

A STUDY OF SUBSTITUTED HYDROXY CHALCONEIMINE IN PREDICTING THE ACOUSTICAL PROPERTIES IN DMSO-WATER MIXTURES IN DIFFERENT CONCENTRATIONS

*Patil N.R.¹ and Thakur S.D.²

¹Department of Chemistry, Govt. V.I.S.H, Amravati, Maharashtra, India, ²Department of Chemistry, R.D.I.K. & N.K.D. College, Badnera (Rly), Amravati, Maharashtra, India

*Author for Correspondence

ABSTRACT

Ultrasonic velocity and density measurements of 2¹-hydroxy – 5¹ – bromo – N – (m - hydroxyphenyl) - chalconeimines (HBN3HPCI) in DMSO – Water mixtures, have been carried out in the concentration range 1×10^{-2} - 4×10^{-2} mol dm⁻³ and in different percentage of dioxane – water mixtures. The experimental data have been used to calculate various acoustical parameters such as velocity, adiabatic compressibility (B_s), apparent molal volume (Φ_v), apparent molal compressibility (Φ_{k(s)}), intermolecular free length (L_f), Specific acoustic impedance (Z_s) and relative association (R_A). The results so obtained are very much comparable and are in good agreement. A molecular interaction study has also been made successfully in the light of these acoustical properties.

Keywords: Acoustical Impedance, Velocity, Mixed Solvents, Substituted Hydroxyl Chalconeimine

INTRODUCTION

Excess – acoustical properties are very helpful in predicting the physico-chemical behaviors and molecular interactions occurring in DMSO-Water mixtures.

Successful attempt has been done made by Thirumaran *et al.*, (2009) to estimate ultrasonic study of intermolecular association through hydrogen bonding in ternary liquid mixtures. Over a wide range of composition which indicates that there exist a strong molecular interaction existing between the unlike molecules.

Bhandakkar *et al.*, (2010) have been studied ultrasonic study of molecular interactions in some bio-liquids. Successful attempts have been made by several workers like Pal *et al.*, (2003) and Aminabhavi *et al.*, (2006) on the measurements and theoretical prediction of various thermo-acoustical properties of liquid binary and ternary mixtures over a wide range of composition and still in progress. In a chemical industry, these properties are very significant in design calculations, heat transfer and mass transfer etc. There has been an increasing interest in the study of molecular interactions and a number of experimental techniques have been used to investigate the interactions in liquid mixtures. Extensive work has been done earlier by Pandey *et al.*, (1999), Pal *et al.*, (1998) and Tamura *et al.*, (2000) to estimate the acoustical properties of liquid mixtures which are helpful in assessing the molecular size and shape, contribution to steric hindrance coupling of torsional oscillations and extent of non-ideal behaviors. The latter, in fact depends on the molecular structure and intermolecular interactions in different combinations. The role of internal pressure in liquid-solution thermodynamics was qualitative but recently its usefulness has been explored for quantitative study of intermolecular forces. Pioneer attempts have been made by Pandey *et al.*, (1993) to show the significance and its correlation with other properties in which the internal pressure has been computed from the knowledge of viscosity (η) density (ρ), and ultrasonic velocity (u) using an indirect method. In the present investigation, ultrasonic velocity (u) and density for substituted chalconeimines have been measured over a wide range of compositions. As for as our knowledge is concerned, no one has attempted in predicting the various acoustical properties for substituted hydroxy chalconeimines so far due to scarcity of experimental data, its theoretical applicability could not be done. Also prediction of solubility parameter of 2¹- hydroxy – 5¹ – bromo – N – (-m- hydroxy phenyl)- chalconeimines for 70% DMSO-water at 307.15 K using experimentally determined value of density, and ultrasonic velocity.

Research Article

Experimental

Component liquid DMSO obtained from the BDH chemical Ltd., Poole, England of Anal R Grade, were purified and dried by usual procedure. Selected substituted hydroxy chalconeimines was: 2¹- hydroxyl - 5¹- Bromo – N – (m-hydroxy phenyl)- chalconeimines. Densities & ultrasonic velocities were measured at determined -307.15 k over a wide range of compositions. Densities were with a bicapillary pycnometer with an estimated error of ±0.001 g/ml. The ultrasonic velocity measurements were made using a crystal controlled variable path ultrasonic interferometer (Mittal Enterprise, Model F-05) of 2 MHz with accuracy of ± 0.03 %.

All the measurements were carried out using a water bath thermo statistically controlled to ± 0.01 K.

Table 1(A): Acoustic parameters of HBN3HPCI (L₅) in DMSO at different percentage of DMSO

% DMSO	Mole fraction of DMSO	Ultrasonic Velocity (u _s) (m/sec)×10 ³	Density (ds) (g/m ³)×10 ⁶	Adiabatic compressibility βs(bar ⁻¹)×10 ⁻¹⁰	Intermolecular free length L _f (A ⁰)×10 ²
100	1	1.5060	1.0537	4.1768	4.1031
90	0.6957	1.5642	1.0568	3.8488	3.9404
80	0.5039	1.6088	1.0660	3.6350	3.8290
75	0.4323	1.6378	1.0735	3.4884	3.7504
70	0.3720	1.6761	1.0786	3.2798	3.6363
60	0.2759	1.7365	1.0844	3.0578	3.5130

Apparent molal volume Φ _v (m ³ /mole)×10 ⁻⁶	Apparent molal Compressibility (Φ _{k(s)})(m ³ mol ⁻¹)×10 ⁻¹⁰	Relative association (R _A)	Specific acoustic impedance Z _s (kg m ⁻² s ⁻¹) ×10 ⁶
66.0759	-32.2729	0.9891	1.5868
-64.2180	-7.1363	0.9948	1.6531
-167.1060	-33.1490	0.9928	1.7151
-26.5200	-34.8854	0.9930	1.7583
-9.6250	-5.4633	0.9912	1.8080
-184.8850	-24.4864	0.9961	1.8831

Theory and Calculation

The apparent molal volumes (φ_v) and apparent molal adiabatic compressibilities φ_{k(s)} in solutions are determined respectively, from density (ds) and adiabatic compressibility (βs) of solution using the equation,

$$\phi_v = (M/ds) + [(d_0 - ds)10^3] / m ds d_0 \quad (1)$$

$$\text{and } \phi_k(s) = [1000 (\beta_s d_0 - \beta_0 ds) / m ds d_0] + (\beta_s M / ds) \quad (2)$$

Research Article

where, d_o is the density of pure solvent, m is molality, M is the molecular weight of solute, β_o is adiabatic compressibility of pure solvent and β_s is adiabatic compressibility of solution.

β_s is calculated from ultrasonic velocity using the equations–

$$\beta_s = 100 / (U_s^2 d_s) \text{ and } \beta_o = 100 / U_o^2 d_o$$

where, U_s is the ultrasonic velocity in the solution in m/s. β_s is in bar^{-1} and $\phi k(s)$ is in $\text{cm}^3 \text{ mol}^{-1} \text{ bar}^{-1}$.

The values of $\phi k(s)$ and ϕv were plotted versus $\sqrt{\text{molality}}$ of solutes. The curves represent the least square and ϕv can be given as–

$$\phi v = \phi^0 v + S_v \sqrt{m} \tag{3}$$

and

$$\phi k(s) = \phi^0 k(s) + S k(s) \sqrt{m} \tag{4}$$

where, $\phi^0 v = v^0$ and $\phi^0 k(s) = k^0$ are the infinite dilution apparent molal volumes and apparent molal adiabatic compressibilities respectively. S_v and $S k(s)$ are the experimental slopes representing ion-ion interactions.

The intermolecular free length (L_f), specific acoustic impedance (z) and relative association (R_A) are calculated by using the following equation–

$$L_f = K \times \sqrt{\beta_s} \tag{5}$$

Where, K is the Jackson's constant.

$$Z = U_s \times d_s \tag{6}$$

$$R_A = \frac{d_s}{d_o} \left(\frac{U_o}{U_s} \right)^{1/3} \tag{7}$$

Table 1(B): Acoustic parameters of HBN3HPCI (L_5) in DMSO at different concentrations of solute in DMSO-Water mixtures

Concentration of ligand (m) (mol/dm ³)	\sqrt{m}	Ultrasonic Velocity (u_s) (m/sec) × 10 ³	Density (ds) (g/m ³) × 10 ³	Adiabatic compressibility β_s (bar ⁻¹) × 10 ⁻¹⁰	Intermolecular free length (L_f) (Å) × 10 ⁻²
0.01	0.1000	1.5365	1.0542	4.0455	4.0378
0.02	0.1414	1.5928	1.0585	3.7341	3.8810
0.03	0.1732	1.6322	1.0687	3.5170	3.7663
0.04	0.2000	1.6513	1.0771	3.4189	3.7125
0.05	0.2236	1.6860	1.0822	3.2602	3.6261

Apparent molal volume Φ_v (m ³ /mole) × 10 ⁻⁶	Apparent molal Compressibility ($\Phi_{k(s)}$) (m ³ mol ⁻¹ bar ⁻¹) × 10 ⁻¹⁰	Relative association (R_A)	Specific acoustic impedance Z_s (kg m ⁻² s ⁻¹) × 10 ⁶
2272.4681	55.0605	0.9964	1.6195
1123.5290	13.1655	0.9891	1.6861
571.385	0.9270	0.9603	1.7447
328.77	-1.9438	0.9937	1.7788
219.7565	-4.5881	0.9922	1.8247

Research Article

RESULTS AND DISCUSSION

In the present investigation different acoustic parameters such as adiabatic compressibility (β_s), apparent molal volume (Φ_v), apparent molal compressibility ($\Phi_{k(s)}$), acoustic impedance (Z_s), relative association (R_A) and intermolecular free length (L_f) of the solutions in different percentage of DMSO and at different concentrations of solute are determined at 307.15 K and presented in table 1(A) & 1(B).

1. It is observed from the Table that the values of β_s decrease with decrease in percentage of DMSO. Also β_s decrease with increase in concentrations in 70% DMSO - water mixture (1×10^{-2})

As concentration increases, number of ions or particles increases in a given volume leading to the decrease in the gap between two species. This indicates that there is a strong interaction between ion and solvent molecules, suggesting a structure promoting behavior of the added solute. This may also imply the decrease in number of free ions, showing the occurrence of ionic association due to strong ion-ion interactions. This fact is again supported from the decrease of β_s with the concentration. When an ion is added to the solvent, it attracts certain solvent molecules towards itself by wrenching the molecule species from the bulk of the solvent. Hence less number of solvent molecules will be made available for the next incoming species. This is known as compression. The decreasing trend of β_s with the concentration may be due to aggregation of solvent molecules around ions supporting the strong ion-solvent interaction.

It could be observed that the values of (Φ_{ks}) are found to be decreasing with increase in the concentration of solute. The negative value of (Φ_{ks}) may be due to loss of compressibility of solute due to strong electrostatic salvations of ions.

Density increases with concentration for DMSO which is due to the shrinkage in the volume which in turn is due to the presence of solute molecules. In other words, the increase in density may be interpreted to the structure former of the solvent due to the added solute.

As stated earlier, due to the presence of solute, there is shrinkage of volume which reflects in the apparent molal volume (Φ_v) which is found to be decreased with increase in the concentration of solute.

It is observed from table 1(b) Φ_v Value are positive at all concentrations of solute, showing that the interactions are insensitive to solvent.

(L_f) increases with increase in percentage of DMSO indicating weak interaction between ion and solvent molecules. This also implies increase in number of free ions showing ionic dissociation but weak ion-ion interactions.

Relative association (R_A) is influenced by two factors:-

i) The breaking up of the solvent molecules on addition of electrolyte to it, and,

(ii) The salvation of ions that is simultaneously present, the former resulting in decrease and later increase of relative association.

Relative association (R_A) decrease initially & increases later due to breaking up of solvent molecules on addition of electrolyte (decrease) and solvation of ion's predominates over the breaking up of the solvent aggregate on addition of solutes, with increasing concentration.

The specific acoustic impedance (Z_s) Values increases with decrease in percentage of DMSO. It also supports weak ion-solvent interactions and electrostatic solvation of ions. Also the acoustic impedance (Z_s) increases with increase in concentration of solute which may be due to lyophobic interaction between solute and solvent molecule.

If decreases linearly with increase in concentration of solute, is due to intermolecular attractive and strong electrostatic forces.

ACKNOWLEDGEMENT

The authors are very much thankful to the principal, R.D.I.K. & N.K.D. Badnera (Rly) Amravati. Also thankful to Govt. V.I.S.H. Amravati for providing necessary facilities. One of the authors (NRP) is thankful to UGC for the award of Rajiv Gandhi National fellowship (RGNF).

Research Article

REFERENCES

- Aminabhavi TM and Bindu G (2006).** Densities, Shear Viscosities, Refractive Indices and Speeds of Sound of Bis (2-methoxy ethyl) Ether + n-alcohols at 298.15, 303.15 and 308.15 K. *Indian Journal of Chemical Technology* **13** 149-153.
- Bhandakkar VD, Chimankar OP and Pawar NR (2010).** Ultrasonic study of molecular interactions in some bio-liquids. *Journal of Chemical and Pharmaceutical Research* **2**(4) 873-877.
- Pal A and Kumar A (1998).** Excess Molar Volumes, Viscosities and Refractive Indices of Diethylene Glycol Dimethyl Ether with Dimethyl Carbonate, Diethyl Carbonate and Propylene Carbonate at (298.15, 308.15 and 318.15) K. *Journal of Chemical Engineering Data* **43** 143-144.
- Pal A, Harish K, Arbad BR and Tekzale AB (2003).** Ultrasonic Speeds And Isentropic Functions Of Mixtures Of N-Methyl -2- Pyrrolidinone And Branched Alcohols At 298.15 K. *Indian Journal of Pure Applied Physics* **41** 113-115.
- Pandey JD, Shukla AK, Tripathi N and Dubey GP (1993).** Internal Pressure, Ultrasonic Velocity and Viscosity of Multi-Component Liquid Systems. *Journal of Physics* **40**(2) 81-84.
- Pandey JD, Vyas VV, Jain JP, Dubey GP, Tripathi N and Dey R (1999).** Speed of Sound, Viscosity and Refractive Index of Multicomponent Systems: Theoretical Predictions from the Properties of Pure Components. *Journal of Molecular Liquids* **81** 123-126.
- Tamura K and Osaki A (2000).** Thermodynamic properties of the binary mixtures of cyclohexanone with globular species. *Thermochimica Acta* 352-354.
- Thirumaran S and George Deepesh (2009).** Ultrasonic study of intermolecular association through hydrogen bonding in ternary liquid mixtures. *Arpan Journal of Engineering and Applied Sciences* **4**(4) 1-11.
- Thirumaran S, Earnest J and Huberd Dhanasundaram B (2010).** Molecular interaction studies on N-Alkanols with DMA in toluene at 303 K. *E- Journal of Chemistry* **7**(2) 465-472.