

## ASSESSMENT OF CARBON SEQUESTRATION OF DURGAWADI SACRED GROVE FROM JUNNAR TEHASIL, PUNE DISTRICT (M.S.)

\*Balasaheb S. Kale<sup>1</sup> and D. D. Ahire<sup>2</sup>

<sup>1</sup>Department of Botany, Loknete Vyankatrao Hiray Arts, Science and Commerce college Panchavati, Nashik-03 M.S (India)

<sup>2</sup>Department of Botany, New Arts, Commerce & Science, College, Ahmednagar

\*Author for Correspondence: E-mail: [kaleunipune@gmail.com](mailto:kaleunipune@gmail.com)

### ABSTRACT

Carbon sequestration is the process of transferring CO<sub>2</sub> from the atmosphere into the soil, plants or plant residues, and other organic solids, which are stored or retained in the unit as part of the soil organic matter. Retention time of sequestered carbon in the soil (terrestrial pool) can range from short-term (not immediately released back to the atmosphere) to long-term (millennia) storage. Western ghats are one of the Mega biodiversity centers in the world and second in India. In the present work, I studied the carbon sequestration potential of the Durgawadi Sacred grove from Junnar Tehasil. It is situated in North-western Ghats in the state of Maharashtra. According to the recorded data from the Durgawadi sacred grove field survey plateau tree cover plants are *Memecylon umbellatum* Burm.f., *Olea dioica* Roxb., *Syzygium cumini* (L.) skeels, *Xantolis tomentosa* (Roxb.) Raf., *Atalantia racemosa* Wight & Arn., *Diospyros montana* Roxb., *Gnidia glauca* (Fres.) Gilg and *Actinodaphne angustifolia* (Bl.) Nees are the most dominant species in terms of carbon sequestration. Durgawadi plateau covers floristic composition present but this plant vegetation is in the form of herb, shrub, tree and these vegetation plant species numbers are 360 in which I am focusing only tree plants were in Durgawadi sacred grove. In this sacred grove above mention plant species were found. These species contribute 106.4 tonnes of biomass per hector for this site. Thus the total biomass contribution by these dominant species in the Durgawadi Sacred grove for the total area of 10.3 ha is 5728.1714 tons.

**Keyword:** Carbon Sequestration, Durgawadi Sacred Grove, Biomass, Northern Western Ghats.


### INTRODUCTION

Carbon sequestration is a biological processes. It involves removing carbon from the atmosphere and depositing it in a reservoir like forest trees. Trees play an important role as carbon sink, during physiological process of photosynthesis, the atmospheric CO<sub>2</sub> is taken by the leaves of trees for the manufacture of food in the form of carbohydrates, proteins, lipids and their combinations thereof like starch, lignin, hemicelluloses; amino acids, proteins etc, and is diverted to other tree parts for disposal and storage (Dey *et al.*, 2010), (Archana *et al.*, 2012) to prevent their release into the environment . Biomass and soil carbon are the two major carbon pools play a major role in living and dead plants (Hamed *et al.*, 2013). The terrestrial forest trees have large accesses to store a sequestered carbon (David *et al.* 2001). Biomass is calculated using density of stems, height of the trees and basal area of the trees in a given location. Therefore, accurate biomass estimation is necessary for better understanding impacts of deforestation on release of carbon compounds responsible for global warming and environmental degradation. Forests sequester and store more carbon than any other terrestrial ecosystem (Jagiwala *et al.*, 2012; Lal *et al.*, 2000; Annissa *et al.*, 2013). The prominent role of evergreen forests in carbon storage and sequestration has increased their importance several fold and brought them to the centre-stage of climate change mitigation strategies (Pandey *et al.*, 2011). Significant amount of biomass and total organic carbon sequestered in standing trees in the campus Dr. B. A.M. University, Aurangabad was

observed by estimation using non destructive method by B. L. Chavan *et al.*, 2010. Carbon sequestration study which is a baseline database, can be used for different research and management purposes by the forest departments (Kale *et al.*, 2010; Ravindranath, Ostwald, 2008). In India two mega biodiverse ecosystems are the Himalayas and Western Ghats, especially the Northern Western Ghats (Savita *et al.*, 2014). Carbon dioxide (CO<sub>2</sub>) is one of the abundant greenhouse gas which causes global warming (Jagruiti Jagiwala *et al.*, 2012). Forests, water bodies and soil are the major sinks accumulating carbon compounds like CO<sub>2</sub> (Archana *et al.*, 2012). The study demonstrates the carbon sequestration potential of ecosystems in Durgawadi Sacred grove.

### **Durgawadi Sacred Grove forest area**

The Durgawadi is one of the largest and florally rich basalt outcrops in Pune District; located 30km from Junnar (INDIA) at the south-west corner in Junnar Taluka between **19°21'96.08"N & 73°64'93.57"E** with an elevation that ranges from **1037–1156 m (MSL)**. It has a semi-evergreen composition of Western ghats which is one of the mega biodiversity center in the world and second in India. It is situated in North-western Ghats in the state of Maharashtra. In the present survey of carbon sequestration potential of Durgawadi Sacred grove from Junnar tehasil; *Memecylon umbellatum* Burm.f., *Olea dioica* Roxb., *Syzygium cumini* (L.) skeels, *Xantolis tomentosa* (Roxb.) Raf., *Atalantia racemosa* Wight & Arn., *Diospyros montana* Roxb., *Gnidia glauca* (Fres.) Gilg and *Actinodaphne angustifolia* (Bl.) Nees are the most dominant species of trees found in this region, in terms of carbon sequestration. Besides this the region is rich in *Rubia cordifolia*, *Sonerila scapigera*, *Habenaria foliosa* and *Cynoglossum amabile*. The vegetation is mainly composed *Memecylon umbellatum* and *Atalantia racemosa* with infrequent occurrence of *Gnidia glauca* and *Actinodaphne angustifolia*. The average plant height is 3m These species contribute 556.13314 tonnes biomass per hectare and the total biomass contribution for the total area of 10.3 ha is 5728.1714 tons.



## **MATERIALS AND METHODS**

**Study area-** Durgawadi Sacred Grove From Junnar Tehasil includes 10.3 hectare, out of that almost 90 % is covered by Semi-Evergreen forest.

**Data Collection-** Extensive and repeated field surveys were carried out from 2015-2016 at both the locations to cover all the seasons of the year. Visit to study area each season and take sample and readings like measure plant height , girth , take soil sample and analysis in laboratory Department of Botany, New Arts, Commerce and Science College, Ahmednagar.

**Methodology (Sampling Technique)-** Trees and lianas were sampled mostly by quadrat method . Both these are area-limited surveys. A line transect of 30m×30m was laid in each vegetation type.

**Biomass carbon** = (above ground biomass carbon + below ground biomass carbon + dead organic matter carbon + Soil carbon) (Hamed *et al.*, 2013).

**Estimation of carbon stock-** Carbon pools are

**Above Ground Biomass was estimated by the method of Chavan *et al.*, (2010)**

1. Size of plot - 30m X 30m for above ground biomass
2. Counting and marking of trees and measurement of each tree for height, GBH (Girth at Breast Height).
3. Density of wood referred from world agroforestry database of wood density.
4. Height of tree is measured using Abney level.
5. Volume of tree trunk calculated from  $= 2 \pi r l$  . (Lal and Singh, 2000)
6. Biomass of tree = volume X density.

**Below Ground Biomass-** Standard root to shoot ratio is used for estimation for underground biomass measurement. A review by Cairns *et al.* (1997), covering more than 160 species from tropical, temperate and boreal forests, estimated a mean root to shoot ratio of 0.26 with a range of 0.18-0.30. **Root Biomass = 0.26 X above ground biomass.** Amount of Carbon present is 50 % of biomass.

**Soil Organic Carbon**

1. Wet digestion or titrimetric determination as per Walkly and Black Method (Walkly, Black, 1934; Tunstall, 2010)
2. Soil bulk density is measured to calculate the volume and total soil carbon.
3. Soil samples were collected at two depths 15cm and 30cm.

**Dead Organic Matter A – Leaf litter and Deadwood**


1. Standing and deadwood biomass calculated as per above ground and below ground biomass.
2. Leaf litter biomass calculated by litter stock change method. leaf litter collected dry weight measured.

## RESULTS

It has been observed that the standing biomass of the above ground wooded parts of the trees and below part like organic soil analysis in Durgawadi Sacred Grove, for the total area of of 10.3 ha was 5728.1714 tons. The average carbon sequestered in Durgawadi Sacred Grove was **556.13314 ton/ha.**

Table no. 1 show quadrates studied at random plant number quadrat . It can be seen that in quadrate no. 07 was having maximum trees i.e 232 and hence maximum carbon of nearly 64 tons/ha. as seen in table no. 02.

Observation table No. 01	
Quadrate	Random Plant Number Wise Quadrat
Q1 -	94
Q2-	90
Q3-	96
Q 4-	119
Q 5-	193
Q 6-	205
Q7 -	232
Q8-	212






Table No. 02: Total carbon tone/ hectare

Total carbon tone/ hectare								
	AGC tone/ha	Leaf Litter c/ha	SOC at 15 cm. tons/ha	SOC at 30 cm. tons/ha	Total C tons/ha	Area (ha)	Carbon tons/ha	Carbon tones/ha
Q 1	1020.86	4.794	3.254	6.585	1035.493	1.7	1760.3381	<b>556.13314</b>
Q 2	173.58	3.948	56.97	5.777	240.275	1.4	336.385	
Q 3	171.43	4.14	0.4805	3.999	180.0495	1.5	270.07425	
Q 4	205.76	3.1812	2.76	56.68	268.3812	1.1	295.21932	
Q 5	383.4	2.5056	3.095	9.738	398.7386	0.9	358.86474	
Q 6	881.83	3.3696	2.534	4.469	892.2026	1.2	1070.64312	
Q 7	650.6	3.384	1.503	63.988	719.475	1.2	863.37	
Q 8	553.93	3.7284	32.25	4.92	594.8284	1.3	773.27692	
						<b>10.3</b>	<b>5728.1714</b>	
<b>Q- Studied Quadrates</b>								
<b>AGC- Above Ground Biomass</b>								
<b>SOC - Soil Organic Carbon at 15 cm. and at 30 cm.</b>								

**CONCLUSIONS**

The future research is to focus on the integration using remotely sensed data. The study is carried out to demonstrate the Carbon Sequestration potential by the Ecosystems in Durgawadi sacred Trees that are good in carbon sequestration can go a long way in mitigating effects of climate change. These studies of Durgawadi Sacred Groves forest area highlight the need for focusing on conservation of forest areas and its ecological value. Random sampling was used to collect different variables since it was

the most versatile and scientific method for estimating above ground biomass. All the terrestrial carbon pools were measured step by step.

## REFERENCES

- Annissa MA, Bam S, Keredin TS, Solomon Raju AJ (2013).** Assessment of biomass and carbon sequestration potentials of standing *Pongamia pinnata* in Andhra University, Visakhapatnam, India. *Bioscience Discovery*, **4**(2) 143-148.
- Archana Warran & Ankur patwardhan (2012).** Carbon Sequestration Potential of Trees in and around Pune City.
- BL Chavan and GB Rasal (2010).** Sequestered standing carbon stock in selective tree species grown in University campus at Aurangabad, Maharashtra, India. *International Journal of Engineering Science and Technology* **2**(7), 3003-3007.
- Brian Tunstall (2010).** Measuring Soil Carbon, Eric.
- David E Reichle and John C Houghton and Robert L Kane (2001).** A review of carbon Sequestration *Science and Technology Opportunities by The U.S. Department of Energy.*
- Dey Seema, Ramachandra TV (2010).** Carbon Sequestration Potential of Ecosystems in Uttara Kannada district, Western Ghats of Karnataka, India. *Lake: Wetlands, Biodiversity and Climate Change.*
- Hamed Haghparast, Azamsadat Delbari and DK Kulkarni (2013).** Carbon sequestration in Pune university campus with special reference to geographical information system (GIS), *Annals of Biological Research*, **4**(4) 169-175.
- Hamed Haghparast, Azamsadat Delbari and D. K. Kulkarni (2013).** Carbon sequestration in Pune university campus with special reference to geographical information system (GIS). *Annals of Biological Research*, **4**(4) 169-175.
- Ian Johnson and Rebecca Coburn (2010).** Trees for carbon sequestration, Primefact 981.
- Jagruti Jagiwala and Dr. Nishith Dharaiya (2012).** Estimating Carbon Sequestration in Forest Vegetation Through Several Scientific Methods: A Review to Find Competent Method For Indian Forests., *Abhinav National Monthly Refereed Journal of Research in Science & Technology*, **2**(3).
- Kale Manish, Shirish P, Ravan A, Roy PS, Singh Sarnam (2009).** Patterns of Carbon Sequestration in Forests of Western Ghats and Study of Applicability of Remote Sensing in Generating CarbonCredits through Afforestation/Reforestation. *Journal of Indian Society for Remote Sensing*. **37** 457–471.
- Lal M and Roma Singh (2000).** Carbon Sequestration Potential of Indian Forests, *Environmental Monitoring and Assessment*, **60** 315–327.
- Rajiv Pandey, GS Rawat and J Kishwan (2011).** Changes in Distribution of Carbon in Various Forest Types of India from 1995-2005, *Silva Lusitana* **19**(1) 41 – 54.
- Ravindranath NH, Ostwald Madelene (2008).** Carbon Inventory Methods: Handbook for greenhouse gas inventory, carbon mitigation and roundwood production projects, Springer used volumetric equations to compute biomass.
- Ravindranath NH, Sudha P, Ramaprasad V, Kulkarni MD, Nagendra MDV (2007).** Development of an agroforestry carbon sequestration project in Khamman district, India. *Mitig adapt strat glob change* **12** 1131-11.
- Savita S Rahangdale & Sanjaykumar R Rahangdale (2014).** Plant species composition on two rock outcrops from the northern Western Ghats, Maharashtra, India. *Journal of Threatened Taxa* **6**(4) 5593–5612,
- The U.S. Department of Agriculture (USDA).** Above-Ground Biomass (Plant) Determinations.
- Walkley A and Black IA (1934).** An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. *Soil Science* **37**, 29-38.