

A COMPREHENSIVE STUDY INTO THE DIVERSITY OF FRESHWATER FISH SPECIES INHABITING LENTIC AND LOTIC AQUATIC ENVIRONMENTS WITHIN THE MATHURA BLOCK, DISTRICT MATHURA, UTTAR PRADESH, INDIA

***Praveen Ojha¹ & Dheeraj Kumar²**

¹Department of Zoology, ²Department of Chemistry,
K.R. (P.G.) College, Mathura, UP, India

*Author for Correspondence: dr.praveenojha11@gmail.com

ABSTRACT

A cumulative total of 31 distinct species has been classified into 6 orders, 23 genera, and 12 families, which includes 4 invasive alien fish species that were systematically collected and identified from both lentic and lotic aquatic habitats within the Mathura block of the District Mathura, Uttar Pradesh. The invasive alien fish species identified encompass *Oreochromis niloticus*, *Cyprinus carpio*, *Hypophthalmichthys molitrix*, and *Clarias gariepinus*; their prevalence is particularly pronounced across extensive sections of the Yamuna River. The investigation indicates that *Clarias gariepinus* represents the sole species of fish consistently found across all designated lentic water bodies within the Mathura block. However, the prevailing circumstances indicate a troubling reduction in the populations of native fish species. Invasive species are progressively securing their establishment as reproductive populations, thereby effectively displacing indigenous fish species.

Keywords: Invasive Alien fishes, Native Fish Fauna, Lentic and Lotic Water Bodies, Yamuna River, Mathura Block

INTRODUCTION

Numerous fish species, many of which are native to India, can be found in both its interior and marine waters (Gopi *et al.*, 2017). There are 5248 genera and 36,640 species of fish in the world. There are 18,614 of them in freshwater (Fricke *et al.*, 2023). In India, there are 3523 species of fish, belonging to 1097 genera, 272 families, and 55 orders. Kosygin *et al.* (2024) offer a comprehensive account of fishes found in India while Van der Laan *et al.* (2023) are used to categorize fishes. In India, more than 300 species of alien fish have been found, including 291 ornamental species, 31 aquaculture species, and 3 larvicidal species.

Mathura, located in the northwest of Uttar Pradesh's Agra division, is situated in the Ganga-Yamuna Doab, inside the Yamuna River basin. The district is located between latitudes 27.14' and 27.58' north and longitudes 77.17' and 78.12' east. Numerous tiny tanks and pokhras can be found all over the district, which is traversed by the Yamuna from north to south. Administratively, Mathura is separated into 735 inhabited villages, 10 blocks, and 3 Tehsils. Unpredictable rainfall and salty water are persistent problems for the district. The local economy also depends heavily on trade, services, tourism, and agriculture. A wide variety of fish species with commercial value can be found in the Yamuna River.

However, the river has been negatively impacted by years of pollution. Fish populations are declining, species composition is changing, and invasive fish species are becoming more prevalent as a result of the water's heavy irrigation use and exposure to large amounts of household and industrial trash. According to the study, the Yamuna River's water is used for drinking, agriculture, and hydropower projects, and water pollution puts local species' habitats at serious risk while unintentionally fostering an environment that is conducive to alien fish. Thus, controlling the introduction of invasive species into the Yamuna

Research Article

River and keeping an eye on the water quality are essential. In both lentic and lotic water basins, such actions could be crucial to the preservation of the ecology and aquatic life.

Due to long-term neglect and lack of upkeep, the majority of the lentic water basins, known as kunds, which were utilized for a variety of functions and were significant freshwater supplies in the Mathura block, have now silted up (Saha *et al.*, 2010). Regarding the fish, no other native species could live there because of the invasive alien fish species *Clarias gariepinus*, also referred to as "Thai magur"

MATERIALS AND METHODS

Study Area: Focus on Mathura Block, Mathura, UP.

Sampling Sites: Selected multiple sites to assess spatial variations in fish diversity

1. **Lotic water body** - Yamuna River stretch viz. 1. Abipur Khadar, 2. Kans Quila, 3. Gokul Barrage 4. Koila Ghat.
2. **Lentic water body** – 1. Shantanu Kund, 2. Talvan, 3. Kumudvan, 4. Krishna Kund 5. Krishna Sarovar.

These water bodies are of perennial and habitat of only single fish species i.e. *Clarias gariepinus* (Thai magur).

Fish Sampling: At different sites and local fish market; identify species using taxonomic keys

Study periods: from year 2022 to 2024.

RESULTS AND DISCUSSION

Results

The entire river's length is used for a variety of human activities, according to the study. The Yamuna River is one of India's most polluted rivers at the moment. The river's deteriorating health is a result of pollution from industrial, agricultural, and residential sources. Practices such as bathing, open defecation, washing clothes, wading cattle, and making religious offerings in public and at home are examples of domestic sources. Practices such as bathing, open defecation, washing clothes, wading cattle, and making religious offerings in public and at home are examples of domestic sources. Industrial discharges introduce heavy metals, phenolics, and other organic contaminants into rivers, negatively affecting water characteristics like temperature, pH, and dissolved oxygen. The Central Pollution Control Board (CPCB) claims that industrial facilities in Uttar Pradesh, Delhi, and Haryana are releasing pollutants into the Yamuna directly.

Fish are frequently seen in Mathura's Kunds, or sacred ponds, and are important to both the local ecology and religious rituals. Because of things like water quality or human activity, some have few or no fish. Fish in kunds are observed for their participation in religious ceremonies, their existence in the natural environment, and the effects of water quality on their populations. According to studies, the diversity and number of native fish can be impacted by elements such as pollution, organic load, metal contamination, and the invasion of exotic fish species.

The data highlights a notable decline in native fauna, primarily due to the following critical factors:

1. Pollution of river and pond water
2. Invasion of alien species into the river and pond ecosystem.
3. Ponds and River is utilized for various human activities.

Discussion

31 freshwater fish species from 6 orders, 23 genera, and 12 families were found during my observations. *Oreochromis niloticus*, *Cyprinus carpio*, *Hypophthalmichthys molitrix*, and *Clarias gariepinus* are the four species were identified as Invasive Alien Fishes (table-1). Fish species were identified from a list and divided into two groups: native and foreign. These invasive species' dominance in biomass over native species is highlighted by their abundance, which can be linked to less stressful environmental conditions.

Research Article

Table 1: List of 31-Species belonging to 6 order, 23 genera and 12 families, of which reported 4-species as Invasive Alien Fishes (mark with*).

Sl. No.	Order	Family	Fish Species Scientific Name
1	Osteoglossiformes	Notopteridae	<i>Notopterus notopterus</i> (Pallas, 1769)
2	Cypriniformes	Danionidae	<i>Cabdio morar</i> (Hamilton, 1822)
3			<i>Amblypharyngodon mola</i> (Hamilton, 1822)
4		Cyprinidae	<i>Catla catla</i> (Hamilton, 1822)
5			<i>Cirrhinus mrigala</i> (Hamilton, 1822)
6			<i>Cirrhinus reba</i> (Hamilton, 1822)
7			<i>Crossocheilus latius</i> (Hamilton, 1822)
8			<i>Cyprinus carpio</i> Linnaeus 1758*
9			<i>Hypophthalmichthys molitrix</i> (Valenciennes, 1844)*
10			<i>Labeo gonius</i> (Hamilton, 1822)
11			<i>Labeo rohita</i> (Hamilton, 1822)
12			<i>Salmostoma bacaila</i> (Hamilton, 1822)
13			<i>Puntius sophore</i> (Hamilton, 1822)
14			<i>Puntius sarana</i> (Hamilton, 1822)
15			<i>Rasbora daniconius</i> (Hamilton, 1822)
16	Siluriformes	Siluridae	<i>Wallago attu</i> (Bloch & Schneider, 1801)
17		Clariidae	<i>Clarias gariepinus</i> (Burchell, 1822)*
18		Heteropneustidae	<i>Heteropneustes fossilis</i> (Bloch, 1794)
19		Bagridae	<i>Mystus vittatus</i> (Bloch, 1794)
20			<i>Mystus cavasius</i> (Hamilton, 1822)
21			<i>Sperata aor</i> (Hamilton, 1822)
22			<i>Mystus seenghala</i> (Sykes)
23		Ailiidae	<i>Eutropiichthys vacha</i> (Hamilton, 1822)
24	Perciformes	Ambassidae	<i>Chanda nama</i> Hamilton, 1822
25			<i>Parambassis ranga</i> (Hamilton, 1822)
26	Cichliformes	Cichlidae	<i>Oreochromis niloticus</i> (Linnaeus, 1758)*
27	Anabantiformes	Anabantidae	<i>Anabas testudineus</i> (Bloch, 1792)
28		Channidae	<i>Channa punctata</i> (Bloch, 1793)
29			<i>Channa striata</i> (Bloch, 1793)
30			<i>Channa gachua</i> (Hamilton, 1822)
31			<i>Channa marulius</i> (Hamilton, 1822)

This circumstance emphasizes even more how deteriorating environmental circumstances are causing native fish populations to gradually decline. According to historical sources, Moza and Mishra (2001), Jhingran (1975), Khan *et al.* (1995), and Mishra *et al.* (2007) reported that there were 70 different species of fish in the Yamuna River. Garg and Saxena (1971) recorded 48 species in the Mathura district, whereas Sharma *et al.* (2014) reported an astounding 93 species in the same river. Numerous researchers have noted the Cyprinidae family's dominance (Bhat, 2003; Sarkar *et al.*, 2009; Vass *et al.*, 2011), and the current analysis confirms this supremacy. Concerns have also been raised about the decline of Indian large carps.

Research Article

According to previous research (Mishra *et al.*, 2007; Vass *et al.*, 2011), the increasing quantity of foreign fish in the Yamuna River supports the idea that these species are gradually establishing breeding populations and displacing the Indian main carps. Reduced water discharge changes both micro- and macro-habitats, favoring non-indigenous species, according to research on alien fish hazards in North American rivers (Kolar and Lodge, 2002). 48 fish species from 13 families have been found in the waterways surrounding Mathura, according to an initial assessment. According to Garg and Saxena (1971), the Cyprinidae family is the most common among them, followed by the Bagridae, Schilbeidae, Clupeidae, and Ophiocephalidae. During the study period, 45 fish species from 8 orders and 16 families were identified from three sampling sites in the Yamuna Rive. Mathura had the highest number of fish species (38), while Delhi had the lowest number (36). The Yamuna River is dominated by cat fishes, as evidenced by the order Cypriniformes having the most species and the family Siluriformes having a total of seven families (Kumar & Saxena, 2021). Ojha (2019) identified 19 species from 16 genera and 9 families, including 4 species that were reported as alien in the Ymuna River. Carnivorous species are the most prevalent in terms of trophic usage, followed by herbivorous and omnivorous species.

4 of these 31 species were designated as invasive alien fishes (table-1). The most prevalent species were those belonging to the Cyprinidae family (12 species), which was followed by the Bagaridae (4 species), Channidae (4 species), Danionidae (2 species), Ambassidae (2 species), Notopteridae (1species), Siluridae (1 species), Claridae (1 species), Heteropneustidae (1 species), Schilbeidae (1 species), Nandidae (1 species), and Anabantidae families (1 species). These invasive species' dominance in biomass over native species is highlighted by their abundance, which can be linked to less stressful environmental conditions. This circumstance emphasizes even more how deteriorating environmental circumstances are causing native fish populations to gradually decline. Due in large part to differences in water quality, a prior research of the Yamuna River revealed 93 fish species in 73 genera, 27 families, and 9 orders, with the most diversity found downstream and the lowest in the midstream (Singh *et al.*, 2014). 112 fish species from 10 orders and 29 families were found in the Yamuna River by Joshi *et al.* (2016). Interestingly, Mathura has the highest chlorine levels (217.4 ppm), a symptom of serious problems with the water quality. While foreign species like *Cyprinus carpio* (common carp) and *Oreochromis niloticus* (tilapia) have increased, particularly in polluted areas, the abundance of Indian main carps, like Labeo and Catla has significantly decreased. Native fish populations have declined as a result of pollution, habitat loss, and the introduction of foreign species.

An overview of the results: Stress the rise of foreign fish species in Mathura Block and the reduction of native fish species.

Conservation Implications: Stress the importance of restoring habitat, reducing pollution, and controlling the introduction of exotic species.

Suggestions

1. Establish routine fish population and water quality monitoring.
2. Encourage community participation and knowledge of conservation initiatives.
3. Implement laws governing the release of exotic species and fishing methods.

ACKNOWLEDGEMENT

For their kind financial support of this research study, I would like to sincerely thank the Department of Higher Education's Research and Development Scheme, Government of Uttar Pradesh. Their financing was essential to this study's successful completion and the achievement of our research objectives.

REFERENCES

Bhat A (2003). Diversity and composition of freshwater fishes in river system of Central Western Ghats, India. *Environmental Biology of Fishes* **68** (1) 25–38.

Research Article

- Fricke R, Eschmeyer WN and Fong JD (2023).** Species by Family/Subfamily. (<http://researcharchive.calacademy.org/research/ichthyology/catalog/SpeciesByFamily.asp>). [Electronic version accessed 10 Oct 2023].
- Garg AK and Saxena MS (1971).** Fishes of Mathura with a key for their identification. *Orissa Veterinary Journal* 6(3/4) 128-132.
- Gopi KC, Mishra SS and Kosygin L (2017).** Pisces. Pp. 527-570. In Current Status of Freshwater Faunal Diversity in India (ed. Kailash Chandra, KC Gopi, D.V. Rao, Valarmathi & JRB Alfred), Zoological Survey of India, Kolkata.
- Jhingran VG (1975).** Fish and fisheries of India, Hindustan Publication, New Delhi.
- Joshi KD, Alam A, Jha DN, Srivastava SK and Kumar V (2016).** Fish diversity, composition and invasion of exotic fishes in river Yamuna under altered water quality conditions. *Indian Journal of Animal Sciences* 86 (8) 957–963.
- Khan MA, Panwar RS, Mathur AK and Rekha (1995).** Final Technical Report on “Investigations on biomonitoring and ecorestoration measures in selected stretches of the rivers-Ganga and Yamuna”, Ganga Project Directorate, Ministry of Environmental and Forests, New Delhi.
- Kolar CS and Lodge DM (2002).** Ecological predictions and risk assessment for alien fishes in North America. *Science* 298 1233–1236.
- Kosygin L, Mohapatra A, Bineesh KK, Sharma I, Jadhav SS and Khynriam D (2024).** Fauna of India Checklist: Pisces. Version 1.0. Zoological Survey India. DOI: <https://doi.org/10.26515/Fauna/1/2023/Choradata:Pisces>.
- Kumar Sumit and Saxena Amita (2021).** Fish composition at three different sites in relation to physico-chemical characteristic of Yamuna River, India. *International Journal of Current Microbiology and Applied Sciences*. 10(1)2203-2216.
doi: <https://doi.org/10.20546/ijcmas.2021.1001.253>.
- Mishra DN, Moza U, Lakra C and Kumar S (2007).** Time scale changes in fisheries of river Yamuna. *Journal of Inland Fisheries Society of India*, 32(2) 48-52.
- Moza U and Mishra DN (2001).** Evaluation of fish biomass and community structure in the context of environmental modifications in upper stretch of river Yamuna. *Applied Fisheries and Aquaculture* 1(3) 17-21.
- Ojha Praveen (2019).** Impact of Invasive fishes on the native fish fauna of River Yamuna at Mathura District, UP. *Global Journal for Research Analysis* 8(6) 142-143.
- Sarkar UK, Gupta BK and Lakra WS (2009).** Biodiversity, ecohydrology, threat status and conservation priority of the freshwater fishes of river Gomti, a tributary of river Ganga (India). *Environmentalist* 30 3-17.
- Sharma AP, Das MK, Samanta S, Pau SK and Bhowmick S (2014).** The Ecology and Fishery Status of River Yamuna. Published by- Director Central Inland Fisheries Research Institute Barrackpore, Kolkata – 700120 West Bengal.
- Saha A, Kansal ML, Mishra GC and Gupta RP (2010).** Restoration of the traditional small water bodies in Braj. *South Asian Journal of Tourism and Heritage* 3 (2) 19-29.
- Singh A, Ansari A, Srivastava S, Verma P and Pathak A (2014).** Impacts of Invasive Fishes on Fishery Dynamics of the Yamuna River, India. *Agricultural Sciences* 5 813-821.
- Van der Laan, Fricke R & Eschmeyer WN (2023).** Eschmeyer's Catalog of Fishes: Classification. (<http://www.calacademy.org/scientists/catalog-of-fishes-classification>). [Electronic version accessed 30 Sep 2023].
- Vass KK, Das MK, Tyagi RK, Katiha PK, Samanta S, Srivastava NP, Bhattacharjya BK, Suresh VR, Pathak V, Chandra G, Debnath D and Gopal B (2011).** Strategies for Sustainable Fisheries in the Indian Part of the Ganga Brahmaputra River Basins. *International Journal of Ecology and Environmental Science* 37(4) 157-218.

Research Article

Copyright: © 2025 by the Authors, published by Centre for Info Bio Technology. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY-NC) license [<https://creativecommons.org/licenses/by-nc/4.0/>], which permit unrestricted use, distribution, and reproduction in any medium, for non-commercial purpose, provided the original work is properly cited.